
Patient Name: Mr. ROHIT KUMAR

Age & Sex : 25 Years / M

Ref. Doctor : Dr. S.K.LUHADIA M.D.(Chest & TB)

Received on : 24/03/2025 Reported on : 25/03/2025

Serial No : **032503240012**



Patient Name: Mr. ROHIT KUMAR

Age & Sex : 25 Years / M

Ref. Doctor : Dr. S.K.LUHADIA M.D.(Chest & TB)

Received on : 24/03/2025 Reported on : 25/03/2025

Serial No : 032503240012

MOLECULAR BIOLOGY (TRUENAT™MTB RIF TEST)

Specimen: SPUTUM

Final Result:- Rif Resistance Not Detected

TruenatTM MTB RIF, Mycobacterium Tuberculosis Rifampicin Resistance Test (for diagnosis of MDR TB), is a follow on test conducted on DNA extracts that have shown MTB positive by TruenatTM MTB.

INTERPRETATION:

RESULT	COMMENTS
RIF SENSITIVE	No mutations detected in the target region of <i>rpoB</i> gene. MDR TB unlikely.
RIF RESISTANT	Mutation detected in the target region of <i>rpoB</i> Gene suggesting MDR TB
INTERMEDIATE	No conclusive result

Test Principle:

The rpoB gene encodes the β subunit of bacterial RNA polymerase. It is the site of mutations that confer resistance to the <u>rifampicin</u> antibacterial agents, such as <u>rifampin</u>. Mutations in rpoB that confer resistance to rifampicin do so by altering residues of the rifampicin binding site on RNA polymerase, thereby reducing binding affinity for rifampicin. Rifampicin resistance is most invariably associated with resistant to isoniazid. Hence, detection of rifampicin resistance is recommended as a reliable proxy for diagnosis of MDR TB.

Pathogen Information:

Tuberculosis (TB) is an infectious disease caused predominantly by the bacillus Mycobacterium tuberculosis. It typically affects the lungs (Pulmonary TB) but can affect other sites as well (Extra pulmonary TB). Pulmonary TB spreads through air and is highly contagious. Over 80% of TB infections are pulmonary and if left untreated, a pulmonary TB patient can infect up to 10-15 other people through close contact over the course of a year.Multidrug-resistant Mycobacterium tuberculosis (MDR-TB) has emerged as a major public health problem worldwide with about 450,000 new cases reported every year.

Target selection:

The target sequence for this test is the RRDR region of the *rpoB* gene (between codon positions 509 and 533), representing mutation hot spots known to be related to rifampic resistance.

Method: Real Time PCR (Melt curve analysis)

Note: Assay result should be interpreted only in the context of other laboratory findings and the total clinical status of patient.

PATHOLOGIST

Patient Name: Mr. ROHIT KUMAR

Age & Sex : 25 Years / M

Ref. Doctor : Dr. S.K.LUHADIA M.D.(Chest & TB)

Received on : 24/03/2025 Reported on : 25/03/2025 Serial No : **032503240012**

